

## T-cell abnormalities in common variable immunodeficiency: the hidden defect

Wong, Gabriel K; Huissoon, Aarnoud P

DOI:

[10.1136/jclinpath-2015-203351](https://doi.org/10.1136/jclinpath-2015-203351)

License:

Creative Commons: Attribution (CC BY)

*Document Version*

Publisher's PDF, also known as Version of record

*Citation for published version (Harvard):*

Wong, GK & Huissoon, AP 2016, 'T-cell abnormalities in common variable immunodeficiency: the hidden defect', *Journal of Clinical Pathology*, vol. 69, no. 8, pp. 672-676. <https://doi.org/10.1136/jclinpath-2015-203351>

[Link to publication on Research at Birmingham portal](#)

### General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

### Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.



## OPEN ACCESS

# T-cell abnormalities in common variable immunodeficiency: the hidden defect

Gabriel K Wong,<sup>1,2</sup> Aarnoud P Huissoon<sup>1,2</sup>

<sup>1</sup>MRC Centre for Immune Regulation, University of Birmingham, Birmingham, UK  
<sup>2</sup>West Midlands Primary Immunodeficiency Centre, Birmingham Heartlands Hospital, Birmingham, UK

**Correspondence to**

Dr Gabriel Wong, MRC Centre for Immune Regulation, University of Birmingham, Medical School, Vincent Drive, Edgbaston, Birmingham B15 2TT, UK; [g.wong@bham.ac.uk](mailto:g.wong@bham.ac.uk)

Received 17 December 2015

Revised 13 March 2016

Accepted 22 March 2016

**ABSTRACT**

This review discusses how the T-cell compartment in common variable immunodeficiency is marked by the premature arrest in thymic output, leading to T-cell exhaustion and immune dysregulation. Although B cells have been the main focus of the disorder, ample experimental data suggest that T-cell abnormalities can be seen in a large proportion of Freiburg Group 1a patients and those suffering from inflammatory complications. The reductions in T-cell receptor excision circles, naïve T cells, invariant NKT cells and regulatory T cells suggest a diminished thymic output, while CD8 T cells are driven towards exhaustion either via an antigen-dependent or an antigen-independent manner. The theoretical risk of anti-T-cell therapies is discussed, highlighting the need for an international effort in generating longitudinal data in addition to better-defined underlying molecular characterisation.

Common variable immunodeficiency (CVID) is a heterogeneous and enigmatic primary immunodeficiency disorder marked by the failure of humoral immunity and immune dysregulation. The discoveries of T-cell-related molecular defects in hyper IgM syndrome and Inducible T-cell costimulatory (ICOS) deficiency highlighted the negative consequences to humoral immunity of the absence of T-cell support,<sup>1 2</sup> and growing experimental data are now suggesting that the T-cell compartment is also disrupted in patients with CVID.

CD4 T-cell lymphopenia was first noted among patients with CVID<sup>3–5</sup> and can dramatically reverse the normal CD4:8 ratio when combined with expansion of senescent CD8 T cells, a feature not usually found in other antibody deficiencies.<sup>6</sup> Despite the array of observed in vitro T-cell abnormalities, the clinical hallmarks of a T-cell immunodeficiency such as recurrent fungal and viral opportunistic infections are lacking in the majority of patients.

The heterogeneity of CVID and the progressive change in its definition over the years make comparing experimental data between studies difficult, but much can still be learnt. This article will examine existing experimental data in the literature regarding the role of T-cell defects in CVID and discuss how infection risk may increase progressively with decreasing thymic output, while the memory compartment shows evidence of abnormal activation. Additionally, we will assess these findings in the context of the revised 2014 European Society for Immunodeficiencies (ESID) diagnostic criteria and discuss their therapeutic implications.

**T-CELLS EXHAUSTION**

In addition to the quantitative alterations in CD4 and CD8 T cells, abnormalities in cytokine production and cell proliferation have been reported in CVID. Experimental data for cytokine output have been well summarised by Varzaneh *et al*,<sup>7</sup> where sluggish production of interleukin (IL) 2 along with deficiencies in other cytokines such as IL-4, IL-5 and IL-10 were reported by a number of studies.<sup>7–9</sup> On the contrary, high levels of inflammatory cytokines such as IL-6, interferon (IFN)- $\gamma$  and tumour necrosis factor (TNF)- $\alpha$  were observed.<sup>7</sup> During the acute phase of a viral infection in immunocompetent individuals, elevations in the classical trio of IL-2, IFN $\gamma$  and TNF $\alpha$  are typically observed. By contrast, studies in chronic hepatitis C and HIV infections demonstrated that T-cell exhaustion, by phenotypic and functional alterations, following a long period of antigenic stimulation was typically marked by the sequential disappearance of IL-2 and then TNF $\alpha$ .<sup>10 11</sup> Therefore, the cytokine signature in CVID mimics an earlier phase of T-cell exhaustion.

Studies of T-cell proliferation also support the idea of a state of T-cell exhaustion. Earlier studies showed that T-cell proliferation to phytohaemagglutinin and OKT3 was normal in patients with CVID but responses may be suboptimal against tetanus toxoid or T-cell receptor (TCR) antibodies.<sup>12</sup> Cyclic AMP-dependent protein kinase A type 1 (PKAI), an inhibitor of T-cell proliferation, can accumulate in exhausted T cells following prolonged antigenic stimulation.<sup>13 14</sup> Using a PKAI selective antagonist, Aukrust *et al*<sup>15</sup> demonstrated that normal proliferation may be restored, suggesting that the poor T-cell proliferation in CVID was secondary to senescence.

Programmed cell death protein 1 (PD-1), an extended family member of CD28 and Cytotoxic T-lymphocyte-associated protein 4 (CTLA-4), functions as an important immune checkpoint for T cells and negatively regulates immune responses. In a murine lymphocytic choriomeningitis virus model, the synergistic use of IL-2 therapy and PD-1 blockade was able to reverse the exhausted immunophenotype of CD8 T cells and to regain control over the infection.<sup>16</sup> Similar to the PKAI antagonist experience, Perreau *et al*<sup>17</sup> showed that in vitro CD4 proliferation in patients with CVID may also be restored by blocking the programmed death ligand 1 and 2 axes (PD-L1/2). While anti-PD-1 antibodies have not been directly tested in CVID, clinical improvement was reported in patients who received IL-2 therapy in the past.<sup>18 19</sup> Along with suboptimal proliferation responses, a small number of studies also suggest an impaired apoptotic function contributing to the accumulation of effete T cells.<sup>20–22</sup>

**To cite:** Wong GK, Huissoon AP. *J Clin Pathol* Published Online First: [please include Day Month Year] doi:10.1136/jclinpath-2015-203351

Overall, both in vivo and in vitro data suggest the presence of potentially reversible T-cell exhaustion in some patients with CVID.

### EXAGGERATED CD8 T-CELL RESPONSES

While the cause of T-cell exhaustion in CVID is not clear, comparable immunophenotypic features can be found with chronic viral reactivation, raising the possibility of an antigenic driver. T cells of patients with CVID, in particular CD8 T cells, exhibit high levels of activation and memory markers including CD29, CD38, CD95, CD45RO and Human leukocyte antigen (HLA)-DR, and low expression of CD27, CD62L and CD45RA.<sup>23–26</sup> Although the expansion of T-cell memory is natural with advancing age, this process is greatly enhanced in CVID. Similarly, a disproportionate increase in terminally differentiated, senescent T cells (CCR7<sup>−</sup>CD45RA<sup>+</sup>) which are positive for CD57 and PD-1 was found across multiple studies, further supporting the presence of a persistent, unregulated cellular immune response in CVID.<sup>6 17 26 27</sup> Unlike T-cell proliferation and cytokine output, immunophenotypic findings are more consistent across studies and affect a larger proportion of patients with inflammatory complications such as polyclonal proliferation, chronic enteropathy, interstitial lung disease and autoimmunity.

To further support the presence of an underlying antigenic driver, CD8 T cells were shown to be oligoclonal by TCR spectratyping.<sup>28–30</sup> As normal repertoire diversity can be seen in age-matched X-linked agammaglobulinaemia patients, this oligoclonality was not thought to be secondary to the deficiency in antibodies.<sup>29</sup> In addition, some patients with CVID were shown to carry very dominant, stable CD8 clonotypes over time.<sup>29</sup> Consistent with the notion of an underlying antigenic driver, a unique set of hyperexpanded TCR $\beta$  complementarity determining region 3 (CDR3) sequences could be found among patients with CVID. Although previous reports highlighted the preferential use of V $\beta$  4, 12 and 17 gene segments in these expanded clonotypes,<sup>20 31</sup> both Ramesh *et al*<sup>32</sup> and our work could not reproduce these results using modern next-generation sequencing approaches and the level of oligoclonality was much more subtle.

Using a pentamer specific for HLA-A2 cytomegalovirus (CMV) peptide (NLV: NLVPMVATV), Marashi *et al* examined the role of CMV as a putative antigenic driver in CVID. A higher frequency (1–1.5%) of CMV NLV-specific CD8 T cells co-expressing high levels of IFN $\gamma$  and TNF $\alpha$  was found in patients with CVID when compared with healthy controls (0.25%), while the frequency of Epstein-Barr virus-specific T cells was not increased.<sup>33</sup> Similarly, Ki76 expression, a proliferation marker, in NLV-specific CD8 T cells was greatly increased in patients with CVID with inflammatory complications (2.4% vs 0.32%).<sup>34</sup> Although causality could not be demonstrated and bystander activation of CMV-specific T cells could not be ruled out, an exaggerated CMV immune response appears to be closely associated with inflammatory complications in CVID.

Autoimmunity is another potential candidate driver for an exaggerated T-cell response. In patients with chronic diarrhoea, T-cell aggregates and nodular lymphoid hyperplasia are often found in intestinal biopsies, mimicking graft-versus-host disease.<sup>35–38</sup> However, detailed examination of tissue T cells is technically challenging and it is currently not possible to differentiate if these histological features are driven by infections, for example norovirus,<sup>39</sup> autoimmunity or other pathological mechanisms. Furthermore, a recent study suggested that tissue

inflammation in CVID is largely driven by CD3<sup>−</sup> innate lymphoid cells as opposed to T cells.<sup>40</sup>

Overall, chronic activation of CD8 T cells in CVID either via an antigen-dependent or antigen-independent manner is likely to contribute to T-cell exhaustion. While CMV appears to be a promising antigenic driver, the role of autoimmunity remains unconfirmed and awaits further study.

### REDUCTION IN THYMIC OUTPUT

Although current studies have not confirmed a putative antigenic driver, the exaggerated T-cell response could also be due to a lack of regulation. By peripheral blood immunophenotyping, Fevang *et al* were the first to demonstrate a lower frequency of CD4<sup>+</sup>CD25<sup>+</sup>FOXP3<sup>+</sup> T cells, an immunophenotype considered characteristic of regulatory T cells (Treg), in patients with CVID. RNA transcript levels for FOXP3 in CD4 T cells were also lower in patients, particularly in those with splenomegaly. To further support this, the frequency of Treg was inversely proportional to neopterin, a serum inflammatory protein.<sup>41</sup> Several studies had since confirmed the reduction in peripheral blood Tregs (CD4<sup>+</sup>CD25<sup>+</sup>FOXP3<sup>+</sup> or CD4<sup>+</sup>CD25<sup>+</sup>CD127<sup>−</sup>) with the greatest deficiencies identified in patients with autoimmune cytopenias or Freiburg Group 1a.<sup>42–45</sup> Carter *et al*<sup>46</sup> also suggested an association between decreased Tregs and CD8 T-cell exhaustion in CVID.

Poor expressions of CTLA-4 and Glucocorticoid-induced TNFR-related protein (GITR) on Tregs were also noted, suggesting a functional deficit.<sup>46 47</sup> Tregs isolated from patients with CVID and autoimmunity had inferior suppressive function when cocultured with autologous CD4<sup>+</sup>CD25<sup>−</sup> T cells, although it is not clear if this phenomenon is primary or secondary, such as thymic sequestration by chronic infections.<sup>42</sup> CTLA-4 haploinsufficiency had been identified in cohorts of patients with CVID and impaired Treg functions.<sup>48 49</sup> An ongoing international collaboration is being carried out to estimate the prevalence of this mutation among patients with CVID.

Altogether, deficiency in Treg provides a logical explanation for the presence of overexpanded and exhausted CD8 T cells, as well as the development of autoimmunity in CVID patients.

The examinations of other thymic derived T cells suggest that the reduction in Tregs may be part of a much broader picture. Invariant NK T cells (iNKT) are a subset of T cells that exhibit both characteristics of NK cells and T cells. They have highly restricted TCRs (V $\alpha$ 24/V $\beta$ 11) and are responsible for a range of immune responses, in particular the control of chronic viral infections.<sup>50</sup> Using CD1d tetramers and V $\alpha$ 24/V $\beta$ 11 antibodies, Fulcher *et al*<sup>51</sup> first demonstrated a significant reduction in iNKT among Freiburg Group 1a CVID patients. The numerical reduction of iNKT was later confirmed by other groups.<sup>52 53</sup> In one study, almost half of the patients had no circulating iNKTs. Stimulation with CD1d tetramer and  $\alpha$ -galcer, a natural ligand for iNKT, also failed to adequately expand the iNKT population.<sup>54</sup> The combined reduction in both Treg and iNKT highlighted a potential problem with thymic output, as both are considered as primary products of the thymus.<sup>55–57</sup> In keeping with this, immunophenotypic data often reveal lower frequencies of naïve (CD45RA<sup>+</sup>CCR7<sup>+</sup>) CD4 and CD8 T cells in CVID. Other T-cell compartments such as central memory, effector memory and terminally differentiated T cells were relatively unaffected or proportionally increased in frequencies, suggesting that T-lymphopenia in CVID is predominantly restricted to the naïve pool in which a numerical deficit was confirmed by bead-calibrated absolute counting. In addition, a reduction in

recent thymic emigrants (CD45RA<sup>+</sup>CD31<sup>+</sup>) was also shown.<sup>6 58</sup> Finally, the T-cell receptor excision circle (TREC) is a gene segment by-product of VDJ rearrangement not replicated during cell division and is considered as the gold standard in measuring thymic output and recent thymic emigrants.<sup>59</sup> As much as a 10× fold reduction in TREC was reported in both CD4 and CD8 T cells of patients with CVID when compared with age-matched healthy controls.<sup>60 61</sup>

Therefore, ample experimental data support the reduction in thymic output in CVID. As proposed by Liston *et al*,<sup>62</sup> reduction in thymic output in other partial T-cell immunodeficiencies may promote autoimmunity by disrupting the balance between effector and regulatory T cells. It is, thus, not surprising that the majority of the findings described so far in this article gravitate towards Freiburg Group 1a patients who have a higher rate of autoimmune complications. The disappearance of iNKT would further hinder B-cell activation and memory formation.<sup>63</sup> It is, however, important to note that many T-cell independent mechanisms such as CLEC16A and Transmembrane activator and calcium modulator and cyclophilin ligand interactor (TACI) polymorphism may also mediate autoimmunity,<sup>64 65</sup> and thymic failure thus represents just one of several mechanisms for patients with CVID to develop autoimmunity. In addition to the increased risk of infection and autoimmunity, T-cell lymphopenia would theoretically impair tumour surveillance, predisposing patients to both haematological and non-haematological malignancies as suggested by Brent and colleagues.<sup>66</sup>

## B-CELL AND T-CELL COLLABORATION

While poor thymic function could lead to the observed T-cell abnormalities in CVID, it offers little explanation for the humoral deficiency. With the exception of ICOS deficiency, the connexion between B-cell and T-cell abnormalities in CVID remains largely speculative.<sup>2</sup> It is possible that the reduction in naïve T cells limits available cognate T cells to support humoral function. In keeping with that, the loss in naïve CD4 T cells was associated with increased infections.<sup>20</sup> IL-21-producing follicular T-helper cells are derived from naïve T cells and play a crucial role in supporting germinal centre functions.<sup>67</sup> However, conflicting evidence exists in the literature regarding their role in the disease process.<sup>6 68</sup>

On the other hand, the combined failure of T cells and B cells warrants further examination of the common progenitors. Interestingly, bone marrow progenitor cells from patients with CVID were less able to form colonies when examined by the methylcellulose assay, indicating a problem even prior to thymic selection.<sup>6 58</sup> Consistent with this hypothesis, Serana *et al*<sup>69</sup> showed that both TREC and Kappa-deleting element recombination circle (KREC) were proportionally reduced in CVID.<sup>69</sup> Compared with severe combined immunodeficiencies such as deficiencies in JAK3, RAG1 and IL7RA, the reduction in TREC and KREC in CVID was only modest and partial.<sup>70</sup> Similar to the excision circle analyses, telomere length of T cells and B cells correlated with each other, further supporting a common progenitor defect driving the simultaneous failure of B cells and T cells.<sup>71</sup> However, telomere shortening was not consistently demonstrated in all studies and more data are required to support or refute this hypothesis.<sup>72</sup>

## DIAGNOSTIC AND THERAPEUTIC CONSIDERATION

Given the prevalence of T-cell abnormalities among patients with CVID, the recent redefinition of CVID by the ESID now emphasises on the use of T-cell immunophenotyping during the diagnostic workup.<sup>73</sup> However, while there is a sound rationale to

exclude patients with recurrent opportunistic infections and late-onset combined immunodeficiency,<sup>74</sup> the arbitrary cut-offs for CD4 lymphopenia and naïve T-cell lymphopenia will exclude a proportion of patients (box 1). Patients with massive splenomegaly and pancytopenia are not particularly susceptible to opportunistic infections and may face diagnostic exclusion. As a significant amount of the T-cell data were generated before the introduction of the revised ESID criteria, it will be of interest to revisit T-cell functions under the new definition in the future.

The success of checkpoint inhibitors such as anti-PD-1 antibodies used in some cancers and experimentally in T-cell exhaustion raises the question if they have a therapeutic role in CVID, with or without recombinant IL-2.<sup>75</sup> However, the clinical application of such therapies is unexplored outside the oncology setting and untested in immunodeficient patients. In addition, there is a real risk of causing or exacerbating the autoimmunity that characterises the very patients who might benefit from such T-cell resuscitation.

Studies of T cells also highlight for the need for specific consideration for immunosuppressants and biologics in CVID. While corticosteroids and rituximab are now widely prescribed for cytopenias and inflammatory conditions in CVID,<sup>76 77</sup> our experiences with anti-T-cell agents are less well established with long-term follow-up data still lacking. Given the diminished thymic output, therapies targeting T cells should be very carefully considered and tailored or they risk compromising the

### Box 1 ESID 2014 revised diagnostic criteria for common variable immunodeficiency<sup>73</sup>

At least one of the following:

- ▶ Increased susceptibility to infection
  - ▶ Autoimmune manifestations
  - ▶ Granulomatous disease
  - ▶ Unexplained polyclonal lymphoproliferation
  - ▶ Affected family member with antibody deficiency
- AND marked decrease of IgG and marked decrease of IgA with or without low IgM levels (measured at least twice; <2SD of the normal levels for their age)

AND at least one of the following:

- ▶ Poor antibody response to vaccines (and/or absent isohaemagglutinins);
- ▶ That is, absence of protective levels despite vaccination where defined
- ▶ Low switched memory B cells (<70% of age-related normal value)

AND secondary causes of hypogammaglobulinaemia have been excluded (see separate list below)

AND diagnosis is established after the 4th year of life (but symptoms may be present before)

**AND no evidence of profound T-cell deficiency, defined as 2 out of the following:**

- ▶ **CD4 numbers/microlitre:** 2–6 years <300, 6–12 years <250, >12 years <200
- ▶ **% naïve CD4:** 2–6 years <25%, 6–16 years <20%, >16 years 10%
- ▶ **T-cell proliferation absent**

T-cell criteria are marked in bold.

AND no evidence of Ataxia telangiectasia (cafe-au lait spots, ataxia, telangiectasia, raised Alpha fetoprotein (AFP)).

ESID, European Society for Immunodeficiencies.



## Review

existing T-cell immunity without replenishment from the thymus in patients with inflammatory complications or Freiburg Group 1a. In addition to improved molecular characterisation, future studies should focus on gathering longitudinal data in this area and test if latent viral control could be affected by immunosuppressive therapies.

Finally, the combined defect in T cells and B cells provokes a consideration for haematopoietic stem cell transplantation (HSCT) in CVID, the only cure for a combined immunodeficiency disorder. A recent multicentre retrospective study involving 25 patients of various clinical presentations, mostly for refractory immune dysregulation and lymphoma, showed that endogenous immunoglobulin production can be revived after a match-related donor or match-unrelated donor bone marrow transplant.<sup>78</sup> However, the procedure carries a significant morbidity and mortality rate (48% overall survival), much exceed that of modern paediatric immunodeficiency HSCT (81–92%).<sup>79</sup> Also, many patients still required immunoglobulin replacement afterwards, limiting its potential application.

## CONCLUSION

Orthogonal evidence from multiple studies indicates an arrest in thymic output coupled with exhaustion in the memory T-cell compartment in patients with CVID and inflammatory complications or Freiburg Group 1a. It is anticipated that the revised ESID diagnostic criteria will result in an increased focus on T-cell abnormalities in CVID and it will be of interest to re-examine many of the observed abnormalities in well-characterised cohorts under the revised criteria. Importantly, this review calls for additional care in the use of immunosuppressants in CVID, with close monitoring of outcomes, perhaps through an international database.

## Take home messages

- Reduction in thymic output is a common feature of CVID.
- T cells are driven towards exhaustion either via an antigen-dependent or an antigen-independent manner.
- Additional care in the use of immunosuppressants is needed to preserve patients' existing immunity.

**Handling editor** Stephen Jolles

**Contributors** GKW and APH wrote the paper.

**Funding** The work was supported by the Wellcome Trust and was completed as part of a clinical training research fellowship for GKW (grant number 099908/Z/12/Z).

**Competing interests** None declared.

**Provenance and peer review** Commissioned; externally peer reviewed.

**Open Access** This is an Open Access article distributed in accordance with the terms of the Creative Commons Attribution (CC BY 4.0) license, which permits others to distribute, remix, adapt and build upon this work, for commercial use, provided the original work is properly cited. See: <http://creativecommons.org/licenses/by/4.0/>

## REFERENCES

- 1 Lee WI, Torgerson TR, Schumacher MJ, *et al.* Molecular analysis of a large cohort of patients with the hyper immunoglobulin M (IgM) syndrome. *Blood* 2005;105:1881–90.
- 2 Grimbacher B, Hutloff A, Schlesier M, *et al.* Homozygous loss of ICOS is associated with adult-onset common variable immunodeficiency. *Nat Immunol* 2003;4:261–8.
- 3 Cunningham-Rundles C, Bodian C. Common variable immunodeficiency: clinical and immunological features of 248 patients. *Clin Immunol* 1999;92:34–48.
- 4 Farrant J, Spickett G, Matamoros N, *et al.* Study of B and T cell phenotypes in blood from patients with common variable immunodeficiency (CVID). *Immunodeficiency* 1994;5:159–69.
- 5 Baumert E, Wolff-Vorbeck G, Schlesier M, *et al.* Immunophenotypical alterations in a subset of patients with common variable immunodeficiency (CVID). *Clin Exp Immunol* 1992;90:25–30.
- 6 Bateman EA, Ayers L, Sadler R, *et al.* T cell phenotypes in patients with common variable immunodeficiency disorders: associations with clinical phenotypes in comparison with other groups with recurrent infections. *Clin Exp Immunol* 2012;170:202–11.
- 7 Varzaneh FN, Keller B, Unger S, *et al.* Cytokines in common variable immunodeficiency as signs of immune dysregulation and potential therapeutic targets—a review of the current knowledge. *J Clin Immunol* 2014;34:524–43.
- 8 Eisenstein EM, Jaffe JS, Strober W. Reduced interleukin-2 (IL-2) production in common variable immunodeficiency is due to a primary abnormality of CD4+ T cell differentiation. *J Clin Immunol* 1993;13:247–58.
- 9 Holm AM, Aukrust P, Aandahl EM, *et al.* Impaired secretion of IL-10 by T cells from patients with common variable immunodeficiency—involve of protein kinase A type I. *J Immunol* 2003;170:5772–7.
- 10 Yi JS, Cox MA, Zajac AJ. T-cell exhaustion: characteristics, causes and conversion. *Immunology* 2010;129:474–81.
- 11 Wherry EJ, Kurachi M. Molecular and cellular insights into T cell exhaustion. *Nat Rev Immunol* 2015;15:486–99.
- 12 Fischer MB, Hauber I, Eggenbauer H, *et al.* A defect in the early phase of T-cell receptor-mediated T-cell activation in patients with common variable immunodeficiency. *Blood* 1994;84:4234–41.
- 13 Ye J, Ma C, Hsueh EC, *et al.* TLR8 signaling enhances tumor immunity by preventing tumor-induced T-cell senescence. *EMBO Mol Med* 2014;6:1294–311.
- 14 Chou JP, Ramirez CM, Ryba DM, *et al.* Prostaglandin E2 promotes features of replicative senescence in chronically activated human CD8+ T cells. *PLoS ONE* 2014;9:e99432.
- 15 Aukrust P, Aandahl EM, Skälhegg BS, *et al.* Increased activation of protein kinase A type I contributes to the T cell deficiency in common variable immunodeficiency. *J Immunol* 1999;162:1178–85.
- 16 West EE, Jin HT, Rasheed AU, *et al.* PD-L1 blockade synergizes with IL-2 therapy in reinvigorating exhausted T cells. *J Clin Invest* 2013;123:2604–15.
- 17 Perreau M, Viganò S, Bellanger F, *et al.* Exhaustion of bacteria-specific CD4 T cells and microbial translocation in common variable immunodeficiency disorders. *J Exp Med* 2014;211:2033–45.
- 18 Cunningham-Rundles C, Bodian C, Ochs HD, *et al.* Long-term low-dose IL-2 enhances immune function in common variable immunodeficiency. *Clin Immunol* 2001;100:181–90.
- 19 Rump JA, Jahreis A, Schlesier M, *et al.* A double-blind, placebo-controlled, crossover therapy study with natural human IL-2 (nhuIL-2) in combination with regular intravenous gammaglobulin (IVIG) infusions in 10 patients with common variable immunodeficiency (CVID). *Clin Exp Immunol* 1997;110:167–73.
- 20 Giovannetti A, Pierdominici M, Mazzetta F, *et al.* Unravelling the complexity of T cell abnormalities in common variable immunodeficiency. *J Immunol* 2007;178:3932–43.
- 21 Lin SJ, Chao HC, Chang KW, *et al.* Effect of interleukin 15 and interleukin 2 on anti-CD3-induced T-cell activation and apoptosis in children with common variable immunodeficiency. *Ann Allergy Asthma Immunol* 2003;91:65–70.
- 22 Goldberg AC, Eliaschewitz FG, Montor WR, *et al.* Exogenous leptin restores in vitro T cell proliferation and cytokine synthesis in patients with common variable immunodeficiency syndrome. *Clin Immunol* 2005;114:147–53.
- 23 Boileau J, Mouillot G, Gérard L, *et al.* Autoimmunity in common variable immunodeficiency: correlation with lymphocyte phenotype in the French DEFI study. *J Autoimmun* 2011;36:25–32.
- 24 Lanio N, Sarmiento E, Gallego A, *et al.* Immunophenotypic profile of T cells in common variable immunodeficiency: is there an association with different clinical findings? *Allergol Immunopathol (Madr)* 2009;37:14–20.
- 25 Viallard JF, Blanco P, André M, *et al.* CD8+HLA-DR+ T lymphocytes are increased in common variable immunodeficiency patients with impaired memory B-cell differentiation. *Clin Immunol* 2006;119:51–8.
- 26 Vlková M, Thon V, Sárkyová M, *et al.* Age dependency and mutual relations in T and B lymphocyte abnormalities in common variable immunodeficiency patients. *Clin Exp Immunol* 2006;143:373–9.
- 27 Mouillot G, Carmagnat M, Gérard L, *et al.* B-cell and T-cell phenotypes in CVID patients correlate with the clinical phenotype of the disease. *J Clin Immunol* 2010;30:746–55.
- 28 Pandolfi F, Trentin L, San Martín JE, *et al.* T cell heterogeneity in patients with common variable immunodeficiency as assessed by abnormalities of T cell subpopulations and T cell receptor gene analysis. *Clin Exp Immunol* 1992;89:198–203.
- 29 Serrano D, Becker K, Cunningham-Rundles C, *et al.* Characterization of the T cell receptor repertoire in patients with common variable immunodeficiency: oligoclonal expansion of CD8(+) T cells. *Clin Immunol* 2000;97:248–58.

- 30 Viillard JF, Ruiz C, Guillet M, *et al.* Perturbations of the CD8(+) T-cell repertoire in CVID patients with complications. *Results Immunol* 2013;3:122–8.
- 31 Duchmann R, Jaffe J, Ehrhardt R, *et al.* Differential usage of T-cell receptor V beta gene families by CD4+ and CD8+ T cells in patients with CD8hi common variable immunodeficiency: evidence of a post-thymic effect. *Immunology* 1996;87:99–107.
- 32 Ramesh M, Hamm D, Simchoni N, *et al.* Clonal and constricted T cell repertoire in Common Variable Immune Deficiency. *Clin Immunol* 2015.
- 33 Marashi SM, Raeiszadeh M, Enright V, *et al.* Influence of cytomegalovirus infection on immune cell phenotypes in patients with common variable immunodeficiency. *J Allergy Clin Immunol* 2012;129:1349–56 e3.
- 34 Marashi SM, Raeiszadeh M, Workman S, *et al.* Inflammation in common variable immunodeficiency is associated with a distinct CD8(+) response to cytomegalovirus. *J Allergy Clin Immunol* 2011;127:1385–93 e4.
- 35 Daniels JA, Lederman HM, Maitra A, *et al.* Gastrointestinal tract pathology in patients with common variable immunodeficiency (CVID): a clinicopathologic study and review. *Am J Surg Pathol* 2007;31:1800–12.
- 36 Ochtrop ML, Goldacker S, May AM, *et al.* T and B lymphocyte abnormalities in bone marrow biopsies of common variable immunodeficiency. *Blood* 2011;118:309–18.
- 37 Washington K, Stenzel Timothy T, Buckley Rebecca H, *et al.* Gastrointestinal pathology in patients with common variable immunodeficiency and X-linked agammaglobulinemia. *Am J Surg Pathol* 1996;20:1240–52.
- 38 Khodadad A, Aghamohammadi A, Parvaneh N, *et al.* Gastrointestinal manifestations in patients with common variable immunodeficiency. *Dig Dis Sci* 2007;52:2977–83.
- 39 Woodward JM, Gkrania-Klotsas E, Cordero-Ng AY, *et al.* The role of chronic norovirus infection in the enteropathy associated with common variable immunodeficiency. *Am J Gastroenterol* 2015;110:320–7.
- 40 Cols M, Rahman A, Maglione PJ, *et al.* Expansion of inflammatory innate lymphoid cells in patients with common variable immune deficiency. *J Allergy Clin Immunol* 2016;137:1206–1215.e6.
- 41 Fevang B, Yndestad A, Sandberg WJ, *et al.* Low numbers of regulatory T cells in common variable immunodeficiency: association with chronic inflammation in vivo. *Clin Exp Immunol* 2007;147:521–5.
- 42 Yu GP, Chiang D, Song SJ, *et al.* Regulatory T cell dysfunction in subjects with common variable immunodeficiency complicated by autoimmune disease. *Clin Immunol* 2009;131:240–53.
- 43 Genre J, Errante PR, Kokron CM, *et al.* Reduced frequency of CD4(+)CD25(HIGH) FOXP3(+) cells and diminished FOXP3 expression in patients with Common Variable Immunodeficiency: a link to autoimmunity? *Clin Immunol* 2009;132:215–21.
- 44 Arumugakani G, Wood PM, Carter CR. Frequency of Treg cells is reduced in CVID patients with autoimmunity and splenomegaly and is associated with expanded CD21lo B lymphocytes. *J Clin Immunol* 2010;30:292–300.
- 45 Horn J, Manguiat A, Berglund LJ, *et al.* Decrease in phenotypic regulatory T cells in subsets of patients with common variable immunodeficiency. *Clin Exp Immunol* 2009;156:446–54.
- 46 Carter CR, Aravind G, Smalle NL, *et al.* CVID patients with autoimmunity have elevated T cell expression of granzyme B and HLA-DR and reduced levels of Treg cells. *J Clin Pathol* 2013;66:146–50.
- 47 Arandi N, Mirshafiey A, Abolhassani H, *et al.* Frequency and expression of inhibitory markers of CD4(+) CD25(+) FOXP3(+) regulatory T cells in patients with common variable immunodeficiency. *Scand J Immunol* 2013;77:405–12.
- 48 Schubert D, Bode C, Kenefeck R, *et al.* Autosomal dominant immune dysregulation syndrome in humans with CTLA4 mutations. *Nat Med* 2014;20:1410–16.
- 49 Kuehn HS, Ouyang W, Lo B, *et al.* Immune dysregulation in human subjects with heterozygous germline mutations in CTLA4. *Science* 2014;345:1623–7.
- 50 Juno JA, Keynan Y, Fowke KR. Invariant NKT cells: regulation and function during viral infection. *PLoS Pathog* 2012;8:e1002838.
- 51 Fulcher DA, Avery DT, Fewings NL, *et al.* Invariant natural killer (iNK) T cell deficiency in patients with common variable immunodeficiency. *Clin Exp Immunol* 2009;157:365–9.
- 52 Carvalho KI, Melo KM, Bruno FR, *et al.* Skewed distribution of circulating activated natural killer T (NKT) cells in patients with common variable immunodeficiency disorders (CVID). *PLoS ONE* 2010;5:e12652.
- 53 Trujillo CM, Muskus C, Arango J, *et al.* Quantitative and functional evaluation of innate immune responses in patients with common variable immunodeficiency. *J Investig Allergol Clin Immunol* 2011;21:207–15.
- 54 Gao Y, Workman S, Gadola S, *et al.* Common variable immunodeficiency is associated with a functional deficiency of invariant natural killer T cells. *J Allergy Clin Immunol* 2014;133:1420–8, 1428.e1.
- 55 Cowan JE, Jenkinson WE, Anderson G. Thymus medulla fosters generation of natural Treg cells, invariant gammadelta T cells, and invariant NKT cells: what we learn from intrathymic migration. *Eur J Immunol* 2015;45: 652–60.
- 56 Nazzari D, Gradolatto A, Truffault F, *et al.* Human thymus medullary epithelial cells promote regulatory T-cell generation by stimulating interleukin-2 production via ICOS ligand. *Cell Death Dis* 2014;5:e1420.
- 57 Berzins SP, Cochrane AD, Pellicci DG, *et al.* Limited correlation between human thymus and blood NKT cell content revealed by an ontogeny study of paired tissue samples. *Eur J Immunol* 2005;35:1399–407.
- 58 Isgrò A, Marziali M, Mezzaroma I, *et al.* Bone marrow clonogenic capability, cytokine production, and thymic output in patients with common variable immunodeficiency. *J Immunol* 2005;174:5074–81.
- 59 Hazenberg MD, Verschuren MC, Hamann D, *et al.* T cell receptor excision circles as markers for recent thymic emigrants: basic aspects, technical approach, and guidelines for interpretation. *J Mol Med* 2001;79:631–40.
- 60 Guazzi V, Aiuti F, Mezzaroma I, *et al.* Assessment of thymic output in common variable immunodeficiency patients by evaluation of T cell receptor excision circles. *Clin Exp Immunol* 2002;129:346–53.
- 61 De Vera MJ, Al-Harhi L, Gewurz AT. Assessing thymopoiesis in patients with common variable immunodeficiency as measured by T-cell receptor excision circles. *Ann Allergy Asthma Immunol* 2004;93:478–84.
- 62 Liston A, Enders A, Sigs OM. Unravelling the association of partial T-cell immunodeficiency and immune dysregulation. *Nat Rev Immunol* 2008;8:545–58.
- 63 Galli G, Pittoni P, Tonti E, *et al.* Invariant NKT cells sustain specific B cell responses and memory. *Proc Natl Acad Sci USA* 2007;104:3984–9.
- 64 Zhang L, Radigan L, Salzer U, *et al.* Transmembrane activator and calcium-modulating cyclophilin ligand interactor mutations in common variable immunodeficiency: clinical and immunologic outcomes in heterozygotes. *J Allergy Clin Immunol* 2007;120:1178–85.
- 65 Li J, Jørgensen SF, Maggadottir SM, *et al.* Association of CLEC16A with human common variable immunodeficiency disorder and role in murine B cells. *Nat Commun* 2015;6:6804.
- 66 Brent J, Guzman D, Bangs C, *et al.* Clinical and laboratory correlates of lung disease and cancer in adults with idiopathic hypogammaglobulinaemia. *Clin Exp Immunol* 2016;184:73–82.
- 67 Craft JE. Follicular helper T cells in immunity and systemic autoimmunity. *Nat Rev Rheumatol* 2012;8:337–47.
- 68 Bossaller L, Burger J, Draeger R, *et al.* ICOS deficiency is associated with a severe reduction of CXCR5+CD4 germinal center Th cells. *J Immunol* 2006;177: 4927–32.
- 69 Serana F, Airò P, Chiarini M, *et al.* Thymic and bone marrow output in patients with common variable immunodeficiency. *J Clin Immunol* 2011;31:540–9.
- 70 Borte S, von Döbeln U, Fasth A, *et al.* Neonatal screening for severe primary immunodeficiency diseases using high-throughput triplex real-time PCR. *Blood* 2012;119:2552–5.
- 71 Visentini M, Cagliuso M, Conti V, *et al.* Telomere-dependent replicative senescence of B and T cells from patients with type 1a common variable immunodeficiency. *Eur J Immunol* 2011;41:854–62.
- 72 Tucker VC, Rye AD, Harrison J, *et al.* Lymphocyte subpopulations from patients with primary antibody deficiency do not show increased telomere erosion. *Immunobiology* 2004;208:455–62.
- 73 Ameratunga R, Brewerton M, Slade C, *et al.* Comparison of diagnostic criteria for common variable immunodeficiency disorder. *Front Immunol* 2014;5:415.
- 74 Malphettes M, Gérard L, Carmagnat M, *et al.* Late-onset combined immune deficiency: a subset of common variable immunodeficiency with severe T cell defect. *Clin Infect Dis* 2009;49:1329–38.
- 75 Philips GK, Atkins M. Therapeutic uses of anti-PD-1 and anti-PD-L1 antibodies. *Int Immunol* 2015;27:39–46.
- 76 Gobert D, Bussell JB, Cunningham-Rundles C, *et al.* Efficacy and safety of rituximab in common variable immunodeficiency-associated immune cytopenias: a retrospective multicentre study on 33 patients. *Br J Haematol* 2011;155:498–508.
- 77 Wong GK, Goldacker S, Winterhalter C, *et al.* Outcomes of splenectomy in patients with common variable immunodeficiency (CVID): a survey of 45 patients. *Clin Exp Immunol* 2013;172:63–72.
- 78 Wehr C, Gennery AR, Lindemans C, *et al.* Multicenter experience in hematopoietic stem cell transplantation for serious complications of common variable immunodeficiency. *J Allergy Clin Immunol* 2015;135:988–97 e6.
- 79 Dvorak CC, Cowan MJ. Hematopoietic stem cell transplantation for primary immunodeficiency disease. *Bone Marrow Transplant* 2008;41:119–26.



# T-cell abnormalities in common variable immunodeficiency: the hidden defect

Gabriel K Wong and Aarnoud P Huissoon

*J Clin Pathol* published online May 6, 2016

---

Updated information and services can be found at:

<http://jcp.bmj.com/content/early/2016/05/06/jclinpath-2015-203351>

---

*These include:*

## References

This article cites 78 articles, 16 of which you can access for free at:  
<http://jcp.bmj.com/content/early/2016/05/06/jclinpath-2015-203351#BIBL>

## Open Access

This is an Open Access article distributed in accordance with the terms of the Creative Commons Attribution (CC BY 4.0) license, which permits others to distribute, remix, adapt and build upon this work, for commercial use, provided the original work is properly cited. See:  
<http://creativecommons.org/licenses/by/4.0/>

## Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

---

## Topic Collections

Articles on similar topics can be found in the following collections

[Open access](#) (99)  
[Immunology \(including allergy\)](#) (1628)

---

## Notes

---

To request permissions go to:

<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:

<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:

<http://group.bmj.com/subscribe/>